

Canadian **U**rological Association
The Voice of Urology in *Canada*



Association des **U**rologues du Canada
La voix de l'urologie au *Canada*

Erectile Dysfunction

Speaker

Trustin Domes MD, MEd, FRCSC

Urologist

Chair, 2021 CUA ED Guideline Committee

Disclosures

- No relationships to disclose.




Learning Objectives


- Describe the physiology of erection
- Classify erectile dysfunction (ED)
- Discuss the assessment of a patient presenting with ED
- Outline the key features of management for a patient presenting with ED



Four Domains of Male Sexual Function



Physiology of Erection: Corporal Level




4 processes:

- 1) Dilation of arteries
- 2) Enlargement of sinusoidal spaces
- 3) Compression of sub-tunical veins
- 4) Increase in cavernosal pressure

Adapted from *BMJ* 1998;316:678–82

Physiology of Erection: Cellular Level



Adapted from *J Sex Med* 2004;1 (3):254-65


Prevalence of ED

ED Rates from Massachusetts Male Aging Study

■ No ED ■ Mild ED ■ Moderate ED ■ Severe ED

n=1290


Age range:
40 to 70 years



J Urol 1994;151:54-61




Severity of ED Increases with Age



Adapted from *J Urol* 1994;151:54-61

Classification of ED



Adapted from *Urology* 1993;42:468-81



Vasculogenic

- Microvascular disease and resulting endothelial dysfunction is a significant underlying cause of ED
- Risk factors for vasculogenic ED:
 - Hypertension
 - Dyslipidemia
 - Diabetes
 - Smoking
 - Pelvic Radiation
 - Vessel injury from trauma/surgery

Some patients may present with ED as their only clinical sign otherwise subclinical vascular disease

ED may be an early marker of cardiovascular disease and may predict future cardiac events, especially in younger men




Am J Med 2014;127:174-82
Am Heart J. 2012;164(1):21-8




Structural

- Conditions that affect the structure and function of the cavernosal bodies
 - Peyronie's Disease is the most common
 - Benign scarring condition of the tunica albuginea of the penis affecting up to 9% of the male population
 - Affects the elastic tissues and expansion capabilities of the corpora cavernosa



Neurogenic

- Numerous neurological conditions can affect erectile function
- CNS:
 - Stroke, Parkinson's Disease, MS, Spinal Cord Injury, Trauma
- PNS:
 - Neuropathies (Diabetes, Alcoholism, Vitamin Deficiencies), Trauma
- Demyelinating diseases:
 - Multiple System Atrophy, MS
- Surgery:
 - Pelvic/Retroperitoneal
 - Most common: Radical Prostatectomy and Proctectomy




Hormonal

- All domains of male sexual function require testosterone
Libido > Mood > Erections > Body Composition Changes
- Causes of Hypogonadism:
 - Idiopathic, often associated with aging (most common)
 - Primary testicular failure: genetic and non-genetic
 - Associated with other medical conditions
 - Liver disease, hemochromatosis, HIV/AIDS, obesity
 - Surgical, traumatic and infectious
 - Hypothalamus/Pituitary
 - Testicle(s)
 - Medications: LHRH agonists and antagonists, opioids, corticosteroids



Psychogenic



ED History

- ED significantly impacts both the patient and their partner
 - Encourage partner involvement in patient assessment
- Onset, duration, consistency and severity of ED
 - Situational versus generalized
 - Response to prior treatments
- Issues with other domains of male sexual function
 - Desire, orgasm, ejaculation, any penile deformity/curvature
- Co-morbid risk factors
 - Medical, surgical, psychological and social factors
- Medication review
 - Antipsychotics (Risperidone and classical), antidepressants (SSRI, SNRI, TCAs) antihypertensives (thiazides, beta-blockers) and anti-androgens



Psychogenic versus Organic ED

Question	Psychogenic ED	Organic ED
Presence of nocturnal erections?	Often present	Reduced
Presence of erection during masturbation or with alternate partners?	Often present	Reduced
Significant recent psychosocial stress?	Strong impact	Minimal impact
Feelings of performance anxiety around sexual activity?	Strong impact	Minimal impact
Situational variability of erectile dysfunction (improved while on vacation)?	Potential for wide variability	Minimal variability

Physical Examination

Area	Factors to be assessed
Overall	Blood pressure, body habitus, virilization, mood, gynecomastia
Penis and groins	Penile length and girth, presence of penile plaques, phimosis, frenular tether, meatal stenosis, quality of femoral pulses
Testicles	Volume and consistency


CUA ED Guideline, 2021



Investigations

- Morning serum testosterone
 - Hypogonadal symptoms (low sexual desire)
 - Failure to respond to PDE5-inhibitors
- Fasting glucose, lipids, hemoglobin A1C
 - Idiopathic or suspected vasculogenic ED cases where occult diabetes and dyslipidemia should be ruled out
- More advanced vascular and neurological testing rarely indicated





Conservative treatment

- Should be discussed with all patients with ED
- Medication review and potential medication change
 - ACE inhibitor/ARB, calcium channel blocker less impact on erections
 - Bupropion, mirtazapine less impact on erections
- Focus on optimizing vascular risk factors and impacts on overall health
- Smoking, alcohol and cannabis negatively impact erectile function
- Increasing physical activity improves erectile function
(conditional recommendation, low certainty of evidence)

Management of Psychogenic ED

- Psychological counseling is typically used in combination with medical management
- Individual and couple counseling sessions are encouraged
- Psychological counseling techniques used by therapists:
 - Anxiety reduction
 - Cognitive-behavioral interventions
 - Discussions regarding adequate sexual stimulation/non-intercourse forms of stimulation
 - Interpersonal assertiveness and couple communication training

Urology Clinics NA 2001;28(2):269-78




Management of ED



Intraurethral alprostadil



Intracavernosal injection




Penile prosthesis



PDE5 inhibitors



Vacuum erection pump device



Least Invasive

Most Invasive

Therapeutic Ladder

Phosphodiesterase Type 5 inhibitors (PDE5-inhibitors)

- Facilitate erection by promoting vascular and cavernosal smooth muscle relaxation in response to sexual stimulation
- Highly effective first-line therapy
 - All three agents in Canada (Sildenafil, Tadalafil, Vardenafil) have similar efficacy with improvement in IIEF-EF score of around 7 in RCTs
- Absolute contraindications: organic nitrate use and allergy
- Side effects:
 - Headache
 - Flushing
 - Dyspepsia
 - Rhinitis
 - Back pain (Tadalafil specific)
 - Alteration in color vision (Sildenafil specific)



Pharmacology of PDE5-inhibitors

Characteristic	Sildenafil	Tadalafil	Vardenafil
Tmax	1.0 h	2.0 h	0.7 h – 0.9 h
Half-life	4 h	17.5 h	4 h – 5 h
Absorption affected by food	Yes	No	Yes
Interaction with alcohol	Yes	Yes	Yes

P. T. 2013; 31:31-5



Daily versus On-demand Tadalafil

- Given the pharmacokinetics of Tadalafil, daily dosing is possible
- Clinical efficacy and side effect profiles are very similar between daily and on-demand regimens
- Decision to prescribe daily versus on-demand PDE5-inhibitor is based on patient preference
- Patient-centered factors influencing daily dosing preference:
 - Increases sexual spontaneity (less concerns regarding timing)
 - Less anticipatory anxiety
 - Treatment of comorbid lower urinary tract symptoms
 - May be more cost-effective depending on frequency of use and whether a low (2.5 mg) or high (5 mg) daily dose is used

CUA ED Guideline, 2021



Low-intensity shockwave therapy (Li-SWT) [experimental]

- Painless shockwaves delivered to different areas of the penis in multiple sessions using a wand-like device
- Hypothesized to improve erectile function through regeneration
 - Induces angiogenesis and nerve regeneration
- Limited research demonstrating clinical efficacy
 - 7 RCTs, varied treatment protocols, 3 trials have high risk of bias
 - IIEF-EF score improvement of 2.07 (95% CL 0.19, 3.96) with moderate certainty of evidence
- NOT Health Canada approved for treatment of ED
- Recommend against Li-SWT as a treatment for patients with ED at this time (conditional, low certainty of evidence)

CUA ED Guideline, 2021



SEXUAL MEDICINE

SOCIETY REPORT

Restorative Therapies for Erectile Dysfunction: Position Statement From the Sexual Medicine Society of North America (SMSNA)



James L. Liu, MD,^{1,*} Kevin Y. Chu, MD,^{2,*} Andrew T. Gabrielson, MD,¹ Run Wang, MD,³ Landon Trost, MD,⁴
Gregory Broderick, MD,⁵ Kelvin Davies, PhD,⁶ Gerald Brock, MD,⁷ John Mulhall, MD,⁸ Ranjith Ramasamy, MD,² and
Trinity J. Bivalacqua, MD, PhD¹



ABSTRACT

Introduction: Current non-invasive treatments for erectile dysfunction (ED) include oral medications, intracavernosal injections, and vacuum-assisted devices. Though these therapies work well for many, a subset of patients have contraindications or are unsatisfied with these options. Restorative therapies for ED are a new frontier of treatments focused on regenerating diseased tissue and providing a potential “cure” for ED.

Aim: The aim of this position statement is to examine existing clinical trial data for restorative therapies and identify elements that require further research before widespread adoption.

Methods: A literature review was performed to identify all clinical trials performed with regenerative therapy for ED. This includes treatments such as stem cell therapy (SCT), platelet rich plasma (PRP), and restorative related technologies like low-intensity shockwave therapy (LiSWT).

Main Outcome Measures: Most clinical trials in restorative therapies were assessed for safety, feasibility, or efficacy. This included recording adverse events, changes in sexual function and erectile function questionnaires, and

Conclusions: Restorative therapies are a promising technology that represents a new frontier of treatment geared towards reversing disease pathology rather than just treating symptoms. However, current published clinical studies are limited. Future work needs to be adequately powered, multi-center, randomized, sham/placebo-controlled trials in well-characterized patient populations to ensure safety and demonstrate efficacy. Until these studies are done, restorative therapies should be reserved for clinical trials and not offered in routine clinical practice. Liu JL, Chu KY, Gabrielson AT, et al. Restorative Therapies for Erectile Dysfunction: Position Statement From the Sexual Medicine Society of North America (SMSNA). *J Sex Med* 2021;9:100343



Second-line therapies

- Vacuum erection pump device
 - Negative pressure promotes blood flow which is trapped by a constriction ring placed at base of penis
 - Good option for patients who cannot tolerate or have contraindications to other medical or surgical options
- Intracavernosal injections (ICI)
 - Injection of vasoactive agent into corpus cavernosum prior to intercourse
 - Highly effective, but risk of pain, bruising, scarring and priapism
- Intraurethral alprostadil (MUSE™)
 - Preferred for patients with needle phobia
 - Risk of urethral/penile pain, very low risk of priapism


Penile prosthesis

- All non-surgical options should at least be discussed with the patient prior to considering surgical intervention
- 2 types of implants
 - Malleable and inflatable
- Very high patient and partner satisfaction rates
- Complications:
 - Infection
 - Device erosion
 - Device mechanical failure




Primary Care


Patient Assessment
Investigations



Health promotion
Modify reversible
causes




First Line
PDE5 Inhibitors
Counseling



Family physician with specialist
interest in men's health
Urology referral

Second Line
Vacuum devices
Intracavernosal injection
Intraurethral therapy



Third Line
Penile prosthesis

Urology referral



Take home points

- ED is common and may be associated with reversible factors or undiagnosed conditions that need to be optimized
- History, physical examination and laboratory investigations (when applicable) narrow down the primary cause of ED in the vast majority of cases
- There are numerous treatment options for ED, typically starting with the least invasive to most invasive using a therapeutic ladder approach
- Family physicians and other primary care providers are the initial point of contact and play a huge role in the diagnosis and treatment of ED
 - Urology referral required only for treatment refractory cases



Canadian **U**rological Association
The Voice of Urology in *Canada*



Association des **U**rologues du Canada
La voix de l'urologie au *Canada*

Erectile Dysfunction

Speaker

Trustin Domes MD, MEd, FRCSC

Urologist

Chair, 2021 CUA ED Guideline Committee